

Summary of research results so far

Kiyoiki Hoshino

18th January, 2025

In analysis for random functions I gave results as follows:

(1) Ogawa integrability ([4])

[6] shows that the Ogawa integral of the Itô process is given with the Itô integral. I and Tetsuya Kazumi showed that the Ogawa integral of the S-type Itô process, which is a noncausal extension of the Itô process, is given with the Skorokhod integral and its adjoint operator.

(2) Identification of random functions from the SFCs ([1, 2, 5])

Let X_t , $t \in [0, L]$ be a random function $X_t = \int_0^t a(t) dB_t + \int_0^t b(t) dt$ driven by a Brownian motion B . To the question whether the coefficients $a(t)$ and $b(t)$ are determined by the stochastic Fourier coefficients (SFCs for short):

$$(e_n, dX) = \int_0^L e_n dX$$

with respect to a CONS $(e_n)_{n \in \mathbb{N}}$ of $L^2([0, L])$ posed in [7], we got the following affirmative answer, employing the SFCs in the case that $\int dB$ is the Skorokhod integral (SFC-Ss in abbr.) and the SFCs in the case that $\int dB$ is the Ogawa integral (SFC-Os in abbr.): here, by FVP, $\mathcal{L}^{r,2}$ we mean the totality of random functions of bounded variation, totality of square itegrable Wiener functionals with differentiability index r , respectively.

- Derivation of random functions from SFC-Ss ([2, 5])

- Derivation of $a(t) \in \mathcal{L}^{1,2}$ and $b(t) \in \mathcal{L}^{0,2}$ ([5]) (extension of the results in [9, 10])

- Derivation of random functions from SFC-Os ([1, 2, 5])

- Derivation of $a(t)$ and $b(t)$ in the case $a(t)$ is written as $a(t) = V_t + M_t + Z_t + W_t$ with $V_t \in \text{FVP}$, an Itô integral process M_t , a Skorokhod integral process Z_t and the Hilbert-Schmidt transform W_t of a functional in $\mathcal{L}^{1,2}$, or a more general random function (extension of the result in [8, 11])

(3) Stochastic differentiability ([3])

The stochastic derivative and quadratic variation is the fundamental operations as the inverse of the stochastic integral. However the sum of random functions X and Y with the quadratic variations does not necessarily have the quadratic variation in general. Then, given a random function V with the quadratic variation, the class:

$$Q(V) = \left\{ X : \text{random function} \left| [X], \frac{dX}{dV} \text{ exists and } \frac{d[X]}{d[V]} = \left| \frac{dX}{dV} \right|^2 \right. \right\}$$

of random functions with the quadratic variation $[]$ and stochastic derivative $\frac{d}{dV}$ with respect to V contains the totality of Itô integral processes, Stratonovich-Fisk integral processes and Skorokhod integral processes and forms a vector space. This result gives a sufficient condition for the sum of random functions X and Y to have the quadratic variation and means that we can unifiedly calculate the stochastic derivative and quadratic variation independent of theory of stochastic integral. Here, the result is used to prove the result in [1] on the identification of random functions from the SFCs mentioned in (2).

References

- [1] K. Hoshino, Identification of random functions from the SFCs defined by the Ogawa integral regarding regular CONSs (Probability Symposium), RIMS Kôkyûroku. **2116**, 95-104, (2019).
- [2] K. Hoshino, Derivation formulas of noncausal finite variation processes from the stochastic Fourier coefficients, Japan Journal of Industrial and Applied Mathematics, vol. 37. **2**, 527-564, (2020).
- [3] K. Hoshino, On the stochastic differentiability of noncausal processes with respect to the process with quadratic variation, Stochastics: An International Journal of Probability and Stochastic Processes, vol. 95. **8**, pp 1446-1473, (2023).
- [4] K. Hoshino, T. Kazumi, On the Ogawa integrability of noncausal Wiener functionals, Stochastics. Vol. 91. **5**, 773-796, (2019).
- [5] K. Hoshino, T. Kazumi, On the Identification of Noncausal Wiener Functionals from the Stochastic Fourier Coefficients, Journal of Theoretical Probability, vol. 32. **4**, 1973-1989, (2019).
- [6] S. Ogawa, The stochastic integral of noncausal type as an extension of the symmetric integrals, Japan J. Appl. Math. **2**, 229-240, (1985).
- [7] S. Ogawa, On a stochastic Fourier transformation, Stochastics. Vol. 85. **2**, 286-294, (2013).
- [8] S. Ogawa, Direct inversion formulas for the natural SFT, Sankhya. The Indian Journal of Statistics, Vol. 80-A, 267-279, (2018).
- [9] S. Ogawa, H. Uemura, On a stochastic Fourier coefficient: case of noncausal function, J.Theoret. Probab. **27**, 370-382, (2014).
- [10] S. Ogawa, H. Uemura, Identification of a noncausal Itô process from the stochastic Fourier coefficients, Bull. Sci. Math. **138**, 147-163, (2014).
- [11] S. Ogawa, H. Uemura, Some aspects of strong inversion formulas of an SFT, Japan Journal of Industrial and Applied Mathematics. 35-1, 373-390, (2018).