

# Lie groups whose moduli spaces of left-invariant Riemannian metrics are one-dimensional

Hiroshi Tamaru (田丸 博士)  
Osaka Metropolitan University / OCAMI

部分多様体幾何とリー群作用 2022  
(東京理科大学)  
2023/03/28

# Abstract

## Framework

- We define  $\mathcal{M}_G$  the moduli space of left-inv. Riem. metrics on a Lie group  $G$ .
- Study the (non)existence of distinguished left-inv. metrics on  $G$ , in terms of  $\mathcal{M}_G$ .

## Problem (not solved)

- Classify  $G$  such that  $\dim \mathcal{M}_G = 1$ .
- Study left-inv. Ricci soliton metrics on such  $G$ , in terms of  $\mathcal{M}_G$ .

## Joint work with

- Masahiro Kawamata
- Yuichiro Taketomi (OMU)

# Background - (1/2)

## Basic Question

- Which homogeneous spaces  $G/K$  admit  $G$ -invariant Einstein or Ricci soliton metrics?

## Def.

- Riem. mfd  $(M, g)$  is **Ricci soliton** if  
 $\exists c \in \mathbb{R}, \exists X \in \mathfrak{X}(M) : \text{Ric}_g = cg + \mathfrak{L}_X g.$

## Note

- A situation of homogeneous Einstein/Ricci solitons depends on the signature of  $c$ .

## Fact (Naber, Petersen-Wylie)

- $(M, g)$  : homogeneous Ricci soliton ( $c > 0$ )  
 $\Rightarrow M \cong [\text{homog. Einstein with } sc > 0] \times [\text{flat}].$

## Fact (cf. Alekseevskii-Kimel'fel'd)

- $(M, g)$  : homogeneous Ricci soliton ( $c = 0$ )  
 $\Rightarrow \text{flat}.$

# Background - (2/2)

## (Generalized) Alekseevskii Conjecture

- $(M, g)$  : homog. Einstein (Ricci soliton),  $c < 0$   
 $\Rightarrow M$  is a solvmanifold?  
(solvable Lie group with left-invariant metric?)

## Fact (Jablonski)

- AC and GAC are equivalent.

## Fact (Böhm-Lafuente, 2023)

- AC is true.

## Motivating Question

- Which solvable Lie groups admit left-invariant Einstein or Ricci soliton metrics?

## Note

- Classification is known only for  $\dim \leq 6$  (Will).
- In higher-dim., the (non-)existence problem is hard to be solved in general.

# Framework - (1/3)

## Recall

- Study the (non-)existence of left-inv. “nice” metrics on Lie groups.

## Difficulty

- For  $n := \dim G$  with Lie algebra  $\mathfrak{g}$ ,  
 $\{\text{left-inv. metrics on } G\} \cong GL(n, \mathbb{R})/O(n)$ ,  
which is too big...
- Not good to ask: nice metric  $\leftrightarrow$  nice point?

## Def. (moduli space)

- $G$  : a simply-conn. Lie group with dim.  $n$ ,
- The **moduli space**  $\mathcal{M}_G$  of left-inv. Riem. metrics on  $G$  is defined by the orbit space of the action of  $\mathbb{R}^\times \text{Aut}(G)$  on  $GL(n, \mathbb{R})/O(n)$ .

## Note

- There is a surjection from  $\mathcal{M}_G$  onto  $\{\text{left-inv. metrics on } G\}/(\text{isometry and scaling})$ .
- It is bijective if  $G$  is nilpotent or completely solvable.

# Framework - (2/3)

## Note

- $\mathcal{M}_G$  is a connected Hausdorff space.
- The points corresponding to principal orbits form a smooth mfd.

## Ex (Hashinaga-T. 2017)

- Consider  $\mathfrak{g} := \text{span}\{e_1, e_2, e_3\}$  with  $a \neq 1$ ,  $[e_1, e_2] = e_2$ ,  $[e_1, e_3] = ae_3$ ,  $[e_2, e_3] = 0$ .
- Then, by a matrix calculation,

$$\mathcal{M}_G \cong \left\{ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \lambda & 1 \end{pmatrix} \mid \lambda \geq 0 \right\} \cong [0, +\infty).$$

- On the corresponding  $G$ , the metric is Ricci soliton if and only if it corresponds to 0.

## Note

- Properties of  $\mathcal{M}_G$  are invariants of  $G$ .
- Question (Expectation): They reflect the properties of left-inv. metrics on  $G$ ?

# Framework - (3/3)

## Note

- The simplest case is  $\dim \mathcal{M}_G = 0$ ,  
i.e.,  $\mathcal{M}_G = \{\text{pt}\}$ ,  
i.e.,  $\mathbb{R}^\times \text{Aut}(G) \curvearrowright GL(n, \mathbb{R})/O(n)$  transitive.

## Thm (Lauret 2003)

- $\dim \mathcal{M}_G = 0$  iff the Lie algebra of  $G$  is  $\mathbb{R}^n$  (abelian),  $\mathfrak{g}_{\mathbb{R}H^n}$  (special type),  $\mathfrak{h}^3 \oplus \mathbb{R}^{n-3}$ .

## Note

- Above three admit left-inv. Ricci solitons; which can be proved without classification.
- Moreover, it satisfies
  - $\mathbb{R}^n$  : flat,
  - $\mathfrak{g}_{\mathbb{R}H^n}$  : negative constant curvature,
  - $\mathfrak{h}^3 \oplus \mathbb{R}^{n-3}$  : non-Einstein.
- Hence, left-inv. Einstein metrics could not be characterized in terms of  $\mathcal{M}_G$ ... but how about Ricci solitons?

# One-dimensional Case - (1/5)

## Conjecture

If  $\dim \mathcal{M}_G = 1$ , then

- $\mathcal{M}_G \cong \mathbb{R}$  or  $[0, +\infty)$ ,
- $\exists$  left-inv. Ricci soliton iff  $\mathcal{M}_G \cong [0, +\infty)$ ,
- a left-inv. Ricci soliton corresponds to 0.

## Note (circumstantial evidence)

- It is true for all examples we know.
- If  $\dim \mathcal{M}_G = 1$ , then we can show that  $G$  is solvable. Therefore left-inv. Ricci soliton is unique (Lauret 2011).

## Prop (a partial answer)

- If  $\mathcal{M}_G \cong [0, +\infty)$ , then 0 corresponds to an isolated orbit.
- A left-inv. metric corresponding to an isolated orbit is Ricci soliton (Taketomi 2022).



# One-dimensional Case - (2/5)

## Note (ongoing idea)

- Recall  $\mathcal{M}_G := \mathbb{R}^\times \text{Aut}(G) \backslash (GL(n, \mathbb{R}) / O(n))$ .
- Consider  $H \curvearrowright M$  : a cohomogeneity one action on an Hadamard mfd.
- If  $H$  is connected, then the orbit space  $H \backslash M$  is  $\mathbb{R}$  or  $[0, +\infty)$  (Berndt-Brück 2001).
- In general  $\mathbb{R}^\times \text{Aut}(G)$  is not connected ...

- We are trying to classify Lie groups with  $\dim \mathcal{M}_G = 1$ .
- Many examples are “almost abelian”.

## Def.

- $\mathfrak{g}$  is **almost abelian** if  $\exists$  codimension one abelian ideal. (Namely  $\mathfrak{g} \cong \mathbb{R} \ltimes \mathbb{R}^{n-1}$ )

# One-dimensional Case - (3/5)

## Fact

- Each  $A \in M(n-1, \mathbb{R})$  gives an almost abelian Lie algebra  $\mathfrak{g}_A = \text{span}\{e_1, \dots, e_n\}$  by

$$\text{ad}_{e_1} = A \quad (\text{on } \text{span}\{e_2, \dots, e_n\}).$$

- $\mathfrak{g}_A \cong \mathfrak{g}_B$  iff  $A$  and  $B$  are similar (conjugate up to nonzero scaling).

## Ex.

- $\mathbb{R}^n$ ,  $\mathfrak{g}_{\mathbb{R}H^n}$ ,  $\mathfrak{h}^3 \oplus \mathbb{R}^{n-3}$  are almost abelian, whose corresponding matrices are

$$O_n, \quad I_n, \quad \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \oplus \mathbb{R}^{n-3}.$$

## Ex.

- Remember  $\mathfrak{g} := \text{span}\{e_1, e_2, e_3\}$  with  $[e_1, e_2] = e_2$ ,  $[e_1, e_3] = ae_3$ ,  $[e_2, e_3] = 0$ .
- This is almost abelian corresponding to

$$\begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix}.$$

# One-dimensional Case - (4/5)

## Thm

$\mathfrak{g}_A$  has one-dim. moduli iff  $A$  is similar to

- $(\alpha I_{n-1}) \oplus (\beta I_1)$  with  $\alpha \neq \beta$ ;

- $I_{n-3} \oplus \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ ;  $\leftarrow M_G \cong \mathbb{R}$

- $\begin{pmatrix} \gamma & 1 \\ -1 & \gamma \end{pmatrix}$  with  $\gamma \geq 0$ ;

- $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \oplus O_{n-5}$ .

$$\left\{ \begin{pmatrix} 1 & 1 \\ \lambda & 1 \end{pmatrix} \mid \lambda > 0 \right\}$$

## Idea of Proof

- $(\Leftarrow)$ : direct calculation.
- $(\Rightarrow)$ : The action is of cohomogeneity one iff the slice representation is very nice iff  $A$  cannot have many eigenvalues...

## Conjecture

- $\dim \mathcal{M}_G = 1$  iff its Lie algebra is one of four almost abelian Lie algebras, or  $\mathfrak{h}^5 \oplus \mathbb{R}^{n-5}$  (5-dim Heisenberg + abelian).

# One-dimensional Case - (5/5)

## Recall

For all known examples with one-dim moduli,

- $\mathcal{M}_G \cong \mathbb{R}$  or  $[0, +\infty)$ ,
- $\exists$  left-inv. Ricci soliton iff  $\mathcal{M}_G \cong [0, +\infty)$ ,
- a left-inv. Ricci soliton corresponds to 0.

$\langle \cdot \rangle$  : Ricci soliton

$\Leftrightarrow \mathbb{R}^* \text{Aut}(G) \cdot \langle \cdot \rangle$  : minimal

$\Leftrightarrow$  " : austere

$\Leftrightarrow$  " : mostly reflective

# Problems

## Conjecture

If  $\mathcal{M}_G$  is one-dimensional, then

- $\mathcal{M}_G \cong \mathbb{R}$  or  $[0, +\infty)$ ,
- $\exists$  left-inv. Ricci soliton iff  $\mathcal{M}_G \cong [0, +\infty)$ ,
- a left-inv. Ricci soliton corresponds to 0.

## Note

- The most hard part would be the non-existence in the case of  $\mathcal{M}_G \cong \mathbb{R}$ ;
- Maybe the classification would be easier...?

## Other Problems

- Study the case of  $\dim \mathcal{M}_G = 2$ .
- Study the case that  $\mathbb{R}^\times \text{Aut}(G) \curvearrowright GL(n, \mathbb{R})/O(n)$  is “nice” (e.g., polar or hyperpolar).
- Study the pseudo-Riemannian version.

$$\mathbb{R}^\times \text{Aut}(G) \curvearrowright GL(p+q, \mathbb{R})/O(p, q)$$

# Ref

- J. Berndt, J.C. Diaz-Ramos, H. Tamaru; Hyperpolar homogeneous foliations on symmetric spaces of noncompact type, *J. Differential Geom.* (2010)
- J. Berndt, H. Tamaru; Homogeneous codimension one foliations on noncompact symmetric spaces, *J. Differential Geom.* (2003)
- J. Berndt, H. Tamaru; Cohomogeneity one actions on symmetric spaces of noncompact type, *J. Reine Angew. Math.* (2013)
- L. P. Castellanos Moscoso, H. Tamaru; A classification of left-invariant symplectic structures on some Lie groups, *Beitrage Algebra Geom.*, to appear.
- T. Hashinaga, H. Tamaru; Three-dimensional solvsolitons and the minimality of the corresponding submanifolds, *Internat. J. Math.* (2017)
- M. Kawamata, Y. Taketomi, H. Tamaru; in progress
- Y. Kondo, H. Tamaru; A classification of left-invariant Lorentzian metrics on some nilpotent Lie groups, *Tohoku Math. J.* (2023)
- Y. Taketomi, H. Tamaru; On the nonexistence of left-invariant Ricci solitons — a conjecture and examples, *Transf. Groups* (2018)

Thank you!